



Okanagan College  
Math 122 (71) Winter 2009  
Term Test One – Marksheets  
Instructor: Clint Lee  
Wednesday, February 4

Student Name: \_\_\_\_\_ Total Marks: \_\_\_\_\_  
40

Problem	Marks	
1 (a)	/3	
1 (b)	/3	
1 (c)	/3	
<b>1 Total</b>		<b>/9</b>
2 (a)	/2	
2 (b)	/3	
2 (c)	/2	
<b>2 Total</b>		<b>/7</b>
<b>3 Total</b>		<b>/4</b>
4 (a)	/2	
4 (b)	/3	
4 (c)	/3	
<b>4 Total</b>		<b>/8</b>
5 (a)	/2	
5 (b)	/3	
<b>5 Total</b>		<b>/5</b>
6 (a)	/2	
6 (b)	/3	
6 (c)	/2	
<b>6 Total</b>		<b>/7</b>
<b>Exam Total</b>		<b>/40</b>





Okanagan College  
Math 122 (71) Winter 2009  
**Term Test One**  
**Instructor: Clint Lee**  
**Wednesday, February 4**

Student Name: \_\_\_\_\_

Total Marks: \_\_\_\_\_  
40

**Instructions.** Do all parts of all 6 questions. Show all work and give explanations where required. You may receive part marks for a question if your work is correct even if the final answer is incorrect. However, if your answer is incorrect and no work or explanation is given, you will receive no marks. The number of points for each question is given in the left margin, total 40. A Formula Sheet is attached. You may use any formula on this sheet on any problem on this test.

1. Evaluate each integral. For any definite integral, give the exact numerical value.

[3] (a)  $\int \left( \frac{2}{x^2} + \frac{3}{2x+1} + e^{x/2} \right) dx$

[3] (b)  $\int \frac{\cos(2 + \ln x)}{x} dx$

[3] (c)  $\int_0^2 \frac{t^3}{\sqrt{4 + 2t^4}} dt$

2. Consider the definite integral

$$\int_2^5 (x^3 - 4x) \, dx$$

- [2] (a) Set up the right endpoint Riemann sum for the definite integral above using  $n$  subintervals,  $R_n$ .  
Do not simplify

- [3] (b) Using Maple the sum in part (a) can be evaluated and simplified to

$$R_n = \frac{63}{4} \frac{(n+1)(7n+3)}{n^2}$$

Use an appropriate limit to determine the exact value of the definite integral above.

- [2] (c) Evaluate the definite integral above using the Fundamental Theorem of Calculus to verify your answer in part (b).

- [4] 3. Consider the definite integral

$$\int_4^7 \frac{x+1}{\sqrt{8-x}} \, dx$$

Evaluate this integral using the substitution  $u = \sqrt{8-x}$ .

4. Wine is being drained from a fermentation vessel in a winery. There is a meter at the outlet from the vessel that measures the rate of flow out of the vessel. The flow rate decreases with time and the rate of decrease is also decreasing. The flow rate  $r(t)$  is measured every 2 minutes over the 16 minutes it takes to drain the tank to the desired level. The measurements, in litres per minute, are given in the table below.

$t, \text{ min}$	0	2	4	6	8	10	12	14	16
$r(t), \text{ L/min}$	52	44	37	31	26	22	19	17	16


- [2] (a) Set up a definite integral for the total amount of wine that is drained out of the fermentation vessel during the 16 minute period.

- [3] (b) Use the Midpoint Rule with  $n$  as large as possible to approximate the value of the definite integral in part (a). Can you determine if this approximation an overestimate or an underestimate? Explain?

- [3] (c) Use the Simpson's Rule with  $n$  as large as possible to approximate the value of the definite integral in part (a). Can you determine if this approximation an overestimate or an underestimate? Explain?

5. The region bounded by the curves  $y = \frac{4}{1+x}$  and  $y = 4 - 2x$  over the interval from  $x = 0$  to  $x = 3$  is shaded in the diagram. It is required to find the area of this region.

- [2] (a) Draw one or more typical approximating rectangles on the diagram and give an expression for the area of each.




- [3] (b) Set up one or more definite integrals that together give the required area. Do not evaluate the integral(s).

6. The graph of a function  $f(t)$  is shown. The curved section of the graph is a circular arc. Define the function  $h(x)$  as


$$h(x) = \int_{-2}^x f(t) dt$$

- [2] (a) Find the values of  $h(0)$ ,  $h(2)$ ,  $h(4)$ , and  $h(6)$ .



- [3] (b) Determine the intervals where the function  $h$  is increasing and decreasing. Identify any critical numbers of the function  $h$  and classify each as a local maximum, local minimum or neither. Give the absolute maximum and minimum values of the function  $h$  on the interval  $[-2, 8]$ .

- [2] (c) Sketch the graph of the function  $h$ . Label the local and absolute extremes identified in part (b) above.



## A Short Table of Integrals

1.  $\int f(g(x))g'(x) dx = \int f(u) du$  where  $u = g(x)$
2.  $\int u dv = vu - \int v du$
3.  $\int u^n du = \frac{1}{n+1}u^{n+1} + C \quad (n \neq -1)$
4.  $\int \frac{du}{u} = \ln|u| + C$
5.  $\int e^u du = e^u + C$
6.  $\int a^u du = \frac{1}{\ln a}a^u + C$
7.  $\int \sin u du = -\cos u + C$
8.  $\int \cos u du = \sin u + C$
9.  $\int \sec^2 u du = \tan u + C$
10.  $\int \sec u \tan u du = \sec u + C$
11.  $\int \csc^2 u du = -\cot u + C$
12.  $\int \csc u \cot u du = -\csc u + C$
13.  $\int \tan u du = \ln|\sec u| + C$
14.  $\int \sec u du = \ln|\sec u + \tan u| + C$
15.  $\int \cot u du = -\ln|\csc u| + C$
16.  $\int \csc u du = -\ln|\csc u + \cot u| + C$
17.  $\int \frac{du}{\sqrt{a^2 - u^2}} = \arcsin\left(\frac{u}{a}\right) + C$
18.  $\int \frac{du}{a^2 + u^2} = \left(\frac{1}{a}\right)\arctan\left(\frac{u}{a}\right) + C$
19.  $\int \sin^n u du = -\frac{1}{n}\sin^{n-1} u \cos u + \frac{n-1}{n} \int \sin^{n-2} u du$
20.  $\int \cos^n u du = \frac{1}{n}\cos^{n-1} u \sin u + \frac{n-1}{n} \int \cos^{n-2} u du$
21.  $\int \tan^n u du = \frac{1}{n-1}\tan^{n-1} u - \int \tan^{n-2} u du$
22.  $\int \sec^n u du = \frac{1}{n-1}\tan u \sec^{n-2} u + \frac{n-2}{n-1} \int \sec^{n-2} u du$
23.  $\int e^{au} \sin bu du = \frac{e^{au}}{a^2 + b^2}(a \sin bu - b \cos bu) + C$
24.  $\int e^{au} \cos bu du = \frac{e^{au}}{a^2 + b^2}(a \cos bu + b \sin bu) + C$

## Short List of Identities

1.  $\cos^2 x + \sin^2 x = 1$
2.  $1 + \tan^2 x = \sec^2 x$
3.  $\cos 2x = \cos^2 x - \sin^2 x = 2\cos^2 x - 1 = 1 - 2\sin^2 x$
4.  $\sin 2x = 2\sin x \cos x$
5.  $\cos^2 x = \frac{1}{2}(1 + \cos 2x)$
6.  $\sin^2 x = \frac{1}{2}(1 - \cos 2x)$

## Numerical Integration Formulas

Midpoint Rule  $M_n = \Delta x \left[ f(\bar{x}_1) + f(\bar{x}_2) + \dots + f(\bar{x}_n) \right]$  where  $\bar{x}_i = \frac{x_{i-1} + x_i}{2}$

Trapezoid Rule  $T_n = \frac{\Delta x}{2} \left[ f(x_0) + 2f(x_1) + \dots + 2f(x_{n-1}) + f(x_n) \right]$

Simpson's Rule  $S_n = \frac{\Delta x}{3} \left[ f(x_0) + 4f(x_1) + 2f(x_3) + \dots + 2f(x_{n-2}) + 4f(x_{n-1}) + f(x_n) \right]$

## Series

The Geometric series is

$$\sum_{i=0}^{n-1} ar^i = a \left( \frac{1 - r^n}{1 - r} \right) = a \left( \frac{r^n - 1}{r - 1} \right)$$

The McLaurin series for the function  $f$  is

$$f(x) = f(0) + f'(0)x + \frac{1}{2!}f''(0)x^2 + \frac{1}{3!}f'''(0)x^3 + \dots + \frac{1}{n!}f^{(n)}(0)x^n + \dots$$

The Taylor series centred at  $x = a$  for the function  $f$  is

$$f(x) = f(a) + f'(a)(x - a) + \frac{1}{2!}f''(a)(x - a)^2 + \frac{1}{3!}f'''(a)(x - a)^3 + \dots + \frac{1}{n!}f^{(n)}(a)(x - a)^n + \dots$$